

Consistency conditions for brane tilings

Ben Davison

10th June 2009

1 Introduction

This paper is concerned with quiver algebras along with superpotential relations arising from brane tilings. In particular, we are interested in the question of calculating Donaldson-Thomas invariants for them, as defined in [7], and with the question of when such algebras can be expected to be Calabi-Yau. The results addressing the first question mark a mild generalisation of the results of [6]. This paper expands upon [6] in two ways: we assume weaker hypotheses, and give proofs for brane tilings on higher genus surfaces, and not just the torus. Moreover, it is hoped that the broad idea which enables us to prove these theorems will find wider application, and lead to a better understanding of the relevant algebras. The observation is that F-term equivalence of paths (the equivalence of paths arising from the superpotential relations) can be represented by a restriction of *homotopy* equivalence of paths. By considering chains of F-term equivalent paths, obtained by sequences of ‘basic equivalences,’ that avoid a tile j when lifted to the universal cover of the surface the brane tiling lives on, we obtain a homotopy of paths that avoids this tile, and so we can use the concept of *winding* to describe this F-term equivalence class. While this paper was being prepared, similar results were obtained independently by Nathan Broomhead in [3].

1.1 Acknowledgements

I would like to thank Balázs Szendrői for introducing me to most of this material, and Kazushi Ueda for his willingness to discuss ideas regarding consistency conditions. I am indebted to Alastair King and Nathan Broomhead for sharing their thoughts, and for suggestions and corrections that led to the rewrite of the final section of this paper.

2 Basic notions

We briefly recall some definitions and fix some notation. We will denote the compact real surface of genus g by M_g . A *brane tiling* of genus g is a tiling F of M_g by convex polygonal tiles and an identification of the 1-skeleton of F with

a bipartite graph, along with a chosen colouring of the vertices by black and white, such that each edge contains one vertex from each coloured set. We will denote this coloured bipartite graph Γ in general. To such data we associate a quiver Q , by the following rules: the underlying graph of Q is the dual graph of F , and the edges are directed so that they go clockwise around the black vertices. Throughout this paper we will talk of edges of the tiling and of the quiver interchangeably - clearly they stand in a 1-1 correspondence. To the quiver Q we associate the usual path algebra $\mathbb{C}Q$. Note that in our notation, if u, v are paths in the path algebra, with u ending at the vertex that v begins at, then uv is the path obtained by first going along u , and then v . Our modules are generally right modules.

We now consider the element $W \in \mathbb{C}Q/[\mathbb{C}Q, \mathbb{C}Q]$ given by taking the sum, over the vertices of Γ , of the cycle of arrows going around the vertex, signed so that those going around the black vertices occur with coefficient $+1$, and those going around the white vertices occur with coefficient -1 . To each edge E of the quiver, there is an associated edge variable $X_E \in \mathbb{C}Q$, and there is a noncommutative derivation

$$\partial/\partial X_E : \mathbb{C}Q/[\mathbb{C}Q, \mathbb{C}Q] \rightarrow \mathbb{C}Q$$

given on each monomial by cyclically permuting each instance of X_E to the front and then deleting it. We define

$$A_F = \mathbb{C}Q/\langle \{(\partial/\partial X_E)W; E \in E(Q)\} \rangle$$

where here and elsewhere $E(Q)$ are the edges of Q and $V(Q)$ are the vertices. A_F^+ is the subalgebra generated by paths of length at least 1. The relations $\{(\partial/\partial X_E)W; E \in E(Q)\}$ will be referred to as the ‘F-terms’, borrowing notation from the physics literature. F-term equivalence of two paths in the path algebra, then, is nothing more than equality modulo the two sided ideal generated by these relations. The relations have form as depicted in Figure (1).

Let $\tilde{M}_g$ denote the universal covering space of M_g . We denote the lift of F (respectively Γ) to $\tilde{M}_g$ by $\tilde{F}$ (respectively $\tilde{\Gamma}$). When we talk of paths in $\tilde{F}$ we mean paths along tiles (precisely, sequences of tiles such that consecutive elements share edges, and such that in the dual quiver the arrow goes the right way across this edge). This is of course the same as a path in the lifted dual quiver (once we have chosen a lift of the starting point of the path), and gives an element of A_F . The tiles of F or $\tilde{F}$ are labelled by the letters i, j, k in this paper. Paths are labelled u, v, w . If u is a path in $\tilde{F}$ or F , $h(u)$ will denote the tile that u ends at, and $t(u)$ will denote the tile that u starts at. If i is a tile in F , the tiles in $\tilde{F}$ that are lifted from i are called i -tiles. Where we wish to distinguish between the tiles of F and the tiles of $\tilde{F}$, we will denote the tiles of F by single letters (e.g. i), and the tiles of $\tilde{F}$ by the single letter corresponding to the tile they project to, along with a subscript (e.g. i_1). Where there is no question of ambiguity, we will use single letters for both, in order to avoid clutter. We will write $[u]$ for the element of A_F that the path u represents.

To avoid any confusion: a letter u outside of square brackets will always be a path along tiles in $\tilde{F}$ or F , and will not be used to denote an element of A_F . In particular, we will only write $u = v$ when u is equal to v as a sequence of tiles, a stronger condition than $[u] = [v]$, which merely implies that u and v are F -term equivalent. We will set $u \sim v$ to mean $[u] = [v]$, i.e. u and v are F -term equivalent.

Definition 2.1. An i -path is a path in $\tilde{F}$ starting from some fixed i -tile. An i -loop is an i -path in $\tilde{F}$ that terminates at an i -tile, i.e. one that projects to a loop from the tile i in M_g . A closed i -loop is an i -loop that terminates at the original i -tile in $\tilde{F}$. A simple i -loop (denoted ω_i) is a closed i -loop that consists of a single revolution around one of the vertices of the chosen i -tile, i.e. the completion of either of the loops in Figure (1). For each arrow a in $\tilde{F}$ there are two F -term equivalent ‘broken loops’ from $h(a)$ to $t(a)$, which we denote ω_a^{br} , when we do not wish to distinguish between them (see Figure (1)). If we do wish to distinguish between them then we write $\omega_{a,B}^{br}$ or $\omega_{a,W}^{br}$, for the broken loop going around the black (respectively white) vertex. If u is a path, u' will denote the path with the last arrow deleted, $'u$ the path with the first arrow deleted.

Remark 2.2. It is easily shown that all simple i -loops are F -term equivalent, and that if u is a path in $\tilde{F}$, $\omega_{t(u)}u \sim u\omega_{h(u)}$. We will frequently use the fact that simple loops can be transported in this way along paths while maintaining F -term equivalence.

Definition 2.3. A path u in the tiling $\tilde{F}$ is minimal if it is not F -term equivalent to any path that contains a simple loop. By Remark (2.2) this is equivalent to the condition that it cannot be written $u \sim v\omega_{h(v)}$ for some v .

We next formalize a generalisation of algebra gradings for quiver algebras. Presumably this idea is standard - it is the natural way to define a ‘grading with many objects’.

Definition 2.4. Let $\mathcal{C}$ be a category, and let A be an algebra over an arbitrary field $\mathbb{K}$. A grading of A by $\mathcal{C}$ is given by a decomposition of the underlying vector space of A

$$A \cong \bigoplus_{\phi \in \text{mor}(\mathcal{C})} V_\phi$$

where we denote the morphisms of $\mathcal{C}$ by $\text{mor}(\mathcal{C})$. This decomposition is required to satisfy the two properties

1. If x, y, z are objects of $\mathcal{C}$, and $\phi \in \text{Hom}(x, y)$ and $\psi \in \text{Hom}(y, z)$, then for all $a \in V_\phi$ and $b \in V_\psi$ $ab \in V_{\psi\phi}$.
2. If x, y, z, w are objects of $\mathcal{C}$, and $y \neq z$, and $\phi \in \text{Hom}(x, y)$ and $\psi \in \text{Hom}(z, w)$, then for all $a \in V_\phi$ and $b \in V_\psi$ $ab = 0$.

This can be generalised to a grading of a category by another category. The canonical example is the grading of $\mathbb{C}Q$ by the unlinearised path category of Q , for an arbitrary quiver Q .

We next come to a definition of consistency. This is equivalent to Condition 4.12 of [6].

Definition 2.5. *Two paths u and v ending at the tile j are weakly equivalent if there is some $n \in \mathbb{Z}_{\geq 0}$ such that $u\omega_j^n \sim v\omega_j^n$. A tiling is consistent if weak equivalence implies F -term equivalence.*

This is essentially a cancellation law. It is equivalent to the condition that for all paths u, v, w , if $[u][w] \neq 0$ and $[u][w] = [v][w]$ then $[u] = [v]$. The set of paths in the quiver (with relations) naturally has the structure of a semigroupoid G_0 , which is in fact a category, if we include zero length paths at the tiles i of F , i.e. u_i , such that $[u_i] = e_i$, where e_i is the idempotent corresponding to the tile i . Precisely:

Definition 2.6. *G_0 is the category having, as objects, the tiles of F . Morphisms in G_0 are given by sequences of tiles $(i_0, i_1, \dots, i_n)$ in F , where $n \geq 0$, up to F -term equivalence, that correspond to paths in the dual quiver Q that respect the orientation of Q . Composition of morphisms is given by composition of paths. The category G_1 is given by formally inverting morphisms of G_0 .*

In [6] it is shown that under certain circumstances A_F possesses a natural torus action that induces a grading. We replace this with the natural grading of A_F by the category G_1 , induced by the natural functor from G_0 to G_1 . The following is a useful technical lemma:

Lemma 2.7. *If u is a closed i -loop in an arbitrary brane tiling F , then there exist $n, m \in \mathbb{Z}_{\geq 0}$ such that $u\omega_i^n \sim \omega_i^m$.*

Proof. Let v be a path along the tiles of $\tilde{F}$, where, for now, we don't assume that v passes through the edges in the correct direction. To such a path we associate a u that does go through every edge in the right direction by the following rule: we go along the edges of v , and each time an edge e passes in the correct direction between the tiles of $\tilde{F}$ we add that edge to u , and each time an edge passes the opposite way, we add one of the broken loops associated to that edge to u . Given such a path v we claim that there is a $n \in \mathbb{N}$ such that, if u is the path associated to v by following the given rule, $u\omega_i^n \sim \omega_i^m$ for some $m \in \mathbb{N}$. This clearly suffices to prove the lemma. Let v be such a path. We can assume that v visits no tile twice, for otherwise we may prove the claim after breaking v into two disjoint loops. Note that adding an edge, and then its inverse, results in adding a simple loop under the rule for associating legal paths to paths that pass the wrong way through some edges. The claim is easy to prove in the case where the number of vertices in the interior of the loop v is 1. But we can easily reduce to this case by adding extra paths along the interior of the loop along with their inverses, and we're done. $\square$

In fact the proof clearly gives us more. As in [6] we can deduce that

Lemma 2.8. *If F is a consistent tiling, and u and v have the same startpoints and endpoints in $\tilde{F}$, they satisfy $u \sim v\omega_{t(v)}^n$ for some $n \in \mathbb{Z}_{n \geq 0}$, or $v \sim u\omega_{t(u)}^n$, for some $n \in \mathbb{Z}_{n \geq 0}$.*

Lemma 2.9. *Let F be a consistent tiling. Let $i, j \in \tilde{F}$ be two tiles. Then there is a minimal path u that starts at i and ends at j .*

Proof. We prove this by induction on $n \geq 0$, the shortest length of a path from i to j . If there is a path of length 0 from i to j we deduce that $i = j$ and so the trivial path of length zero, starting at i , is a minimal path from i to j . So assume $i \neq j$, and, using the inductive hypothesis, assume there is a $k \in \tilde{F}$ such that there is an arrow a from k to j in $\tilde{Q}$, the lift of the dual quiver for F , and a minimal path u starting at i and ending at k . Assume also that ua is not minimal, for else we deduce immediately that there is a minimal path from i to j . It follows that $ua \sim v\omega_j$ for some path v . We can pick ω_j to go through k , since all simple loops are F-term equivalent. This gives us a decomposition $ua \sim v\omega_a^{br}a$. By consistency we deduce that $u \sim v\omega_a^{br}$. Now from the definition of minimality of paths we deduce that a sub-path of a minimal path is minimal, from which we obtain that v is a minimal path to j . $\square$

Remark 2.10. *If F is a consistent tiling then, using Lemma (2.8) and Remark (2.2), we deduce that a path u in $\tilde{F}$ is minimal if and only if it is not F-term equivalent to a path that visits some tile twice.*

The consistency condition can be restated as saying that the natural functor from G_0 to G_1 is faithful. There is a natural grading of the algebra A_F by the groupoid G_1 . Finally, it is seen that weak equivalence of two paths u and v is equivalent to $[u]$ and $[v]$ having the same G_1 -grade, and consistency is equivalent to each G_1 -graded piece of A_F being 1-dimensional. We will make use of the following:

Definition/Proposition 2.11. *Let F be a consistent brane tiling. For i a fixed tile in $\tilde{F}$, we define the partial ordering on tiles in $\tilde{F}$, denoted $<_i$, by the rule: $j <_i k$ if there is a minimal path from i to k passing through j .*

We prove that this is well-defined:

1. Transitivity: Let i, j, k, l be tiles of $\tilde{F}$, satisfying $j <_i k$ and $k <_i l$. Then there is a minimal path u from i to l passing through k . We decompose u as $u = vw$ where v is a path from i to k , and w is a path from k to l . Now v is minimal, for otherwise u certainly isn't. From Lemma (2.8) we deduce that all minimal paths from i to k are F-term equivalent, so since $j <_i k$ we see that v is F-term equivalent to a path passing through j . It follows that u is F-term equivalent to a path passing through j .
2. Antisymmetry: Let i, j, k be tiles of $\tilde{F}$. Assume for a contradiction that $j <_i k$ and $k <_i j$. Then there is a minimal path u from i to k passing




Figure 1: A basic F-term equivalence takes us from one broken loop to the other.

through j . We decompose this path as $u = vw$ where v is a path from i to j and w is a path from j to k . As in the check of transitivity, we deduce that v is minimal. Since we have $k <_i j$, we deduce, as before, that v is F-term equivalent to a path passing through k . It follows that u is F-term equivalent to a path that visits the tile k twice, and is not minimal, by Remark (2.10).

3. Reflexivity: Let i, j be tiles of F . Let u be a minimal path from i to j , which exists by Lemma (2.9). Then this path demonstrates that $j \leq_i j$.

3 F-term equivalence and homotopy equivalence

We introduce here a vital heuristic for the proofs in this paper. Consider again the terms coming from the noncommutative derivations of the superpotential. They give equivalences between the ‘broken loops’ as depicted in Figure (1). In fact the relations are given precisely by identifying each set of broken loops. We say there is a *basic* F-term equivalence between two paths u and v if

$$\{u, v\} = \{w_1 \omega_{a,B}^{br} w_2, w_1 \omega_{a,W}^{br} w_2\}$$

for some paths w_1, w_2 in F , and some arrow a in F .

Let γ be a (suitably well behaved) continuous path through M_g , in the sense of a continuous function

$$\gamma : [0, 1] \rightarrow M_g.$$

As long as γ passes edges of F the ‘right’ way (i.e. with the black vertex to the right), we can naturally associate to it a path (in the sense used throughout the rest of this paper) along the dual quiver Q . In this way we obtain a function

$$\Psi : Cont_F([0, 1], \tilde{M}_g) \rightarrow paths_F$$

where $Cont_F([0, 1], \tilde{M}_g)$ is just the set of continuous paths such that the above construction gives a well defined path along F , crossing each edge of $\tilde{F}$ with the right orientation, and $paths_F$ is the set of paths of F . Note that this function is surjective.

If $u \sim v$ then there is a chain of basic F-term equivalences taking u to w . Let γ_u and γ_w be two continuous paths in $Cont_F([0, 1], \tilde{M}_g)$ such that

$$\begin{aligned}\Psi(\gamma_u) &= u \\ \Psi(\gamma_w) &= w.\end{aligned}$$

Say v is obtained from u by applying a basic F-term equivalence. Then there exist paths u_1 and u_2 in $\tilde{F}$ such that $u = u_1 \omega_{a,B}^{br} u_2$ and $v = u_1 \omega_{a,W}^{br} u_2$. There exist $a, b \in [0, 1]$ such that the copy of $\omega_{a,B}^{br}$ indicated in the given decomposition of u is obtained by

$$\omega_{a,B}^{br} = \Psi(\gamma_u|_{[a,b]} \circ s)$$

where s is just a scaling function $s(x) = x(b - a) + a$. As indicated in Figure (1) there is a homotopy

$$H : [0, 1] \times [0, 1] \rightarrow \tilde{M}_g$$

satisfying $H|_0 = \gamma_u|_{[a,b]} \circ s$ and $H|_1 = \nu$, where

$$\nu : [0, 1] \rightarrow \tilde{M}_g.$$

is a new path satisfying $\Psi(\nu) = \omega_{a,W}^{br}$. We pick this homotopy to be invariant along the line $[0, 1] \times \{0\}$, and invariant along the line $[0, 1] \times \{1\}$. This is just the same as stipulating that the homotopy fixes the endpoints, so we can construct a new homotopy H' given by

$$\begin{aligned}H'(t, x) &= \gamma_u(x) && \text{if } t \notin [a, b] \\ &= H(t, s^{-1}(x)) && \text{otherwise.}\end{aligned}$$

This gives a homotopy such that $\Psi(H'|_0) = u$ and $\Psi(H'|_1) = v$. Note that, as indicated in the figure, we can pick H' so that $image(H' : [0, 1]^2 \rightarrow \tilde{M}_g)$ does not intersect any of the tiles not included in u or v .

The crucial observation is that if there is a chain of basic F-term equivalent paths between two paths u and w through $\tilde{F}$, none of which go through some tile k of $\tilde{F}$, then we can produce $\gamma_w, \gamma_u \in Cont_F([0, 1], \tilde{M}_g)$ such that $\Psi(\gamma_w) = w$, $\Psi(\gamma_u) = u$ and an endpoint-preserving homotopy H from γ_w to γ_u such that

$$Image(H : [0, 1]^2 \rightarrow \tilde{M}_g) \cap k = \emptyset.$$

This motivates the following definition.

Definition 3.1. A pair of F-term equivalent paths (u, v) is k -unwound if there is a sequence of paths $u = u_0, \dots, u_n = v$ such that u_s is related to u_{s+1} by a basic F-term equivalence, for every s , and none of the u_s pass through tile k .

In particular, if $u, v \in \text{paths}_F$ are a pair of k -unwound paths then we can find a pair of paths $\gamma_u, \gamma_v \in \text{Cont}([0, 1], \tilde{M}_g)$ that avoid k , and such that $\Psi(\gamma_u) = u$ and $\Psi(\gamma_v) = v$, and such that there is an endpoint preserving homotopy between γ_u and γ_v , considered as paths in $\tilde{M}_g \setminus k$. While this definition makes sense for tilings on all universal covers of oriented surfaces $\tilde{M}_g$, for $g = 0$ it is not useful, since all paths between two chosen tiles in $\tilde{M}_g \setminus k$ are homotopy equivalent. We could of course define the k -winding number of the ordered pair (u, v) , or the $H_1(\tilde{M}_g \setminus K)$ -class of the ordered pair (u, v) for K any union of tiles of $\tilde{F}$, and u, v two paths avoiding K . However, we have no need for these added complications here (except perhaps for Remark (8.2)).

4 3-Calabi-Yau property

We proceed to give a detailed proof of the following theorem, in which a winding argument plays a key part.

Theorem 4.1. *Let $A := A_F$ be the algebra associated to a consistent tiling F on a surface M_g for $g \geq 1$. For each i a tile of F there is a G_1 -graded projective resolution of*

$$S_i := e_i A / e_i A^+$$

given by

$$0 \longrightarrow [\omega_i]A \xrightarrow{\tau} \bigoplus_{a \in E_1} [\omega_a^{br}]A \xrightarrow{\sigma} \bigoplus_{b \in E_2} [b]A \xrightarrow{\rho} e_i A \quad (1)$$

where E_1 is the set of edges going into i , and E_2 is the set of edges going out of i . The maps are defined as follows: ρ is the sum of the obvious inclusion maps. Let $a \in E_1$. Then there are natural inclusions $\iota_1 : [\omega_a^{br}]A \rightarrow [b_1]A$ and $\iota_2 : [\omega_a^{br}]A \rightarrow [b_2]A$, where b_1 is the first arrow in $\omega_{a,B}^{br}$ and b_2 is the first arrow in $\omega_{a,W}^{br}$. We let σ be the direct sum of $\iota_1 - \iota_2$ for $a \in E_1$. Finally, there are natural inclusions $[\omega_i]A \rightarrow [\omega_a^{br}]A$, for each $a \in E_1$, and we let τ be the sum of these.

Proof. That (1) is a chain complex follows straight from the definitions of the maps. All the maps in (1) are clearly G_0 -graded and therefore G_1 -graded. There is a morphism f of complexes from (1) to the complex consisting of zeroes in every position except for a copy of S_i in the zeroeth position, induced by the map

$$e_i A \rightarrow S_i.$$

We must check that this induces isomorphisms on homology, i.e. that $H^i(f)$ is an isomorphism for all i .

$H^0(f)$ is an isomorphism, since the kernel of the map from $e_i A$ to S_i is generated by paths of length 1, all of which lie in the image of ρ . Now we need just to check that the complex (1) is exact away from the zeroeth position.

Injectivity of τ follows from the injectivity of each of its components. Now restrict attention to the G_1 -graded piece of this chain complex corresponding to $[u] \in A$ where u is a path from a chosen i -tile to some j -tile. This path is fixed for the rest of the proof. Note that for each of the summands $[v]A$, in each of the modules of (1), consistency implies that $\dim([v]A)_{([u])} = 1$ if u is F-term equivalent to some path starting with v , and $\dim([v]A)_{([u])} = 0$ otherwise.

Define $E'_2 \subset E_2$ to be the subset of E_2 consisting of those arrows a such that there is a path v with $v \sim u$ such that v passes through $h(a)$. Note that the arrow a is the unique minimal path F-term equivalent to a , since we cannot apply any F-term equivalences to it. Say $a \in E'_2$. Then there is some minimal path v , F-term equivalent to u , such that v passes through $h(a)$. We can decompose v as $v = v_1 v_2$ where v_1 is a path from $t(a)$ to $h(a)$, and v_2 is a path from $h(a)$ to $h(u)$. Using the fact that sub-paths of minimal paths are minimal, we deduce that $v_1 = a$. We deduce that $a \in E'_2$ if and only if there is a path v , F-term equivalent to u , such that v starts with a .

We define E'_1 to be the subset of E_1 consisting of those a such that there is a path v such that $v \sim u$ and v passes through $t(a)$. Again, by the fact that broken loops are the unique minimal paths F-term equivalent to broken loops, we deduce that $a \in E'_1$ if and only if there is some v such that $v \sim u$ and v begins with ω_a^{br} .

Say, to start with, that u is not a minimal path. It follows that u is F-term equivalent to some path starting with a loop ω_i . It follows that $E'_1 = E_1$ and $E'_2 = E_2$. We deduce that the $[u]$ -graded piece of the complex is given by

$$0 \longrightarrow \mathbb{C} \xrightarrow{\tau_{[u]}} \bigoplus_{a \in E_1} \mathbb{C}_a \xrightarrow{\sigma_{[u]}} \bigoplus_{b \in E_2} \mathbb{C}_b \xrightarrow{\rho_{[u]}} \mathbb{C} \quad (2)$$

where $\mathbb{C}_a \cong \mathbb{C}$ for each $a \in E_1 \cup E_2$, $\tau_{[u]}$ is the diagonal map, $\rho_{[u]}$ takes vectors to the sum of their entries, and the map $\sigma_{[u]}$ takes 1_a to $1_{b_1} - 1_{b_2}$, where b_1 is the edge bordering a at a black vertex, and b_2 is the edge bordering a at a white vertex. Exactness of (2) is clear.

So assume that u is minimal. We start by investigating what $E'_1 \cup E'_2$ looks like (considered as edges in $\tilde{\Gamma}$, rather than arrows in the dual quiver, for now). We say a path w demonstrates that $a \in E'_1$ if $w \sim u$ and w starts with ω_a^{br} . Similarly, a path w demonstrates that $a \in E'_2$ if it starts with a , and $w \sim u$. Trivially, if a path demonstrates that an edge is in E'_1 , it demonstrates that a neighbouring edge is in E'_2 . Given two edges e and f in $E'_1 \cup E'_2$, we consider two paths w_e and w_f demonstrating that e and f , respectively, belong to $E'_1 \cup E'_2$. Since w_e and w_f are F-term equivalent there is a sequence of paths $w_e = w_1, \dots, w_t = w_f$ such that each w_s is related to its predecessor by a basic F-term equivalence. Each of these paths demonstrates that one or two connected members of $E_1 \cup E_2$ is a member of $E'_1 \cup E'_2$ (two in the case that w_s starts with a broken loop). The set of members of $E_1 \cup E_2$ that w_s demonstrates is in $E'_1 \cup E'_2$ is connected to the set that w_{s-1} demonstrates is in $E'_1 \cup E'_2$. We deduce that $E'_1 \cup E'_2$ is connected. Also, since $\omega_{a,B}^{br} \sim \omega_{a,W}^{br}$, we deduce that if $e \in E'_1$, the two edges in E_2 connected to e are in E'_2 . We deduce that either

- (A) the set of edges in $E'_1 \cup E'_2$ forms an unbroken line, with edges from E'_2 at either end, or
- (B) these edges form an unbroken loop, i.e. $E'_1 \cup E'_2 = E_1 \cup E_2$.

Restricting to the $[u]$ -grade of the complex, it becomes

$$0 \longrightarrow \bigoplus_{a \in E'_1} \mathbb{C} a \xrightarrow{\sigma_{[u]}} \bigoplus_{b \in E'_2} \mathbb{C} b \xrightarrow{\rho_{[u]}} \mathbb{C} \quad (3)$$

where $\rho_{[u]}$ is still the summation map, and we modify the description of $\sigma_{[u]}$ by the prescription that $1_b = 0$ if $b \notin E'_2$. The zero at the first place is the result of our picking u to be minimal, which is equivalent to $([\omega_i]A)_{[u]} = 0$. From this chain complex and our description of $E'_1 \cup E'_2$ it is clear that exactness of (3) is equivalent to situation (B) not occurring.

So say we are in situation (B) for our grade $[u]$. We construct a sequence of basic F-term equivalences winding around tile i . Let a, b, c be edges of tile i as indicated in Figure (2). Since we are in situation (B), there is a path u_1 , as indicated in Figure (2) which is F-term equivalent to u , and starts with $\omega_{a,B}^{br}$. Again, by the supposition that $E'_1 \cup E'_2 = E_1 \cup E_2$, there is a path u_2 , F-term equivalent to u , starting with $\omega_{c,W}^{br}$. Both u_1 and u_2 start with the arrow b , and by consistency we have $'u_1 \sim' u_2$ (recall that for a path v we denote by $'v$ the path obtained by deleting the first arrow of v). Furthermore, $'u_1$ and $'u_2$ are i -unwound, by minimality.

There is a basic F-term equivalence that replaces $\omega_{c,W}^{br}$ with $\omega_{c,B}^{br}$, taking u_2 to a new path u_3 . We next repeat the reasoning of the above paragraph, winding anticlockwise around the tile i .

We proceed in this way around tile i until we get to a new path u_∞ starting with $\omega_{a,B}^{br}$, F-term equivalent to u . We note that none of the paths in this chain of F-term equivalent paths passes through tile i , except to start at it, by minimality of u . Again, by minimality of u , and cancellation for consistent tilings, $'u_\infty$ and $'u_1$ are F-term equivalent and i -unwound, but this is absurd, since we have constructed u_∞ to wind around i . $\square$

The following definition is taken from [4].


Definition 4.2. *An algebra A is Calabi-Yau of dimension d provided there is an A -bimodule quasi-isomorphism*

$$f : A \longrightarrow A^! [d]$$

such that $f = f^! [d]$, where for an arbitrary A -bimodule the functor $M \mapsto M^!$ is given by $\mathbf{RHom}_{A \otimes A^{op} - \text{mod}}(M, A \otimes_{\mathbb{C}} A)$.

Theorem 4.3. *Let A_F be the algebra associated to a consistent brane tiling F with dual quiver Q . The complex in (1) is a projective resolution of S_i , for each i , if and only if A_F is a Calabi-Yau algebra of dimension 3.*

Figure 2: Winding around tile i_0 .



Proof. The method of proof is a modified version of the argument of [6].

Define the algebra

$$R := \bigoplus_{i \in V(Q)} \mathbb{C},$$

the direct sum of $|V(Q)|$ copies of the algebra $\mathbb{C}$. As a vector space this is generated by elements 1_i for $i \in V(Q)$. Then A_F is naturally a R bimodule - there is an algebra morphism $\iota : R \rightarrow A_F$ sending 1_i to e_i . If S is an R -bimodule, S defines an endofunctor for the category of left R -modules via the tensor product over R . We say that S is (left) flat if this functor takes exact sequences to exact sequences, and is (left) faithfully flat if the exactness of $S \otimes M^\bullet$ for some complex $M^\bullet$ of left R -modules implies the exactness of $M^\bullet$. The following facts are easy to verify.

1. All R -bimodules are (left) flat.
2. An R -bimodule S is (left) faithfully flat if and only if $S \cdot 1_i \neq 0$ for all i .

To each edge $a \in E(Q)$ we associate an R -bimodule E_a in the natural way: E_a has underlying vector space $\mathbb{C}$, and

$$\begin{aligned} 1_k \cdot m \cdot 1_l &= m && \text{if } k=t(a), l=h(a) \\ &= 0 && \text{otherwise.} \end{aligned}$$

We pick a generator of the underlying vector space of E_a and denote it 1_a . We define

$$E := \bigoplus_{a \in E(Q)} E_a.$$

There is a natural basis for the underlying vector space of E given by the elements 1_a , for $a \in E(Q)$.

R is a sub-algebra of A_F and inherits a natural G_1 -grading. Furthermore we give E a G_1 -grading by setting the grade of 1_a to be the grade of $[a]$, considered as an element of A_F .

Next, to each edge $a \in E(Q)$ we associate an R -bimodule E_a^* . This has underlying vector space $\mathbb{C}$, and

$$\begin{aligned} 1_k \cdot m \cdot 1_l &= m \quad \text{if } k=h(a), l=t(a) \\ &= 0 \quad \text{otherwise.} \end{aligned}$$

We define

$$E^* := \bigoplus_{a \in E(Q)} E_a^*.$$

There is a natural basis for the underlying vector space of E^* given by elements denoted 1_a^* , where $1_a^* \in E_a^*$. This R -bimodule has a G_1 -grading given by setting the grade of 1_a^* to be the grade $[\omega_a^{br}]$.

Finally we define a graded R -bimodule L . As an *ungraded* R -bimodule this is isomorphic to R . We shift the grading by setting the grade of the copy of 1_i in L , for some vertex $i \in V(Q)$, to be the G_1 -grade $[\omega_i]$.

We must now show that the sequence

$$0 \longrightarrow A \otimes_R L \otimes_R A \xrightarrow{j} A \otimes_R E^* \otimes_R A \longrightarrow A \otimes_R E \otimes_R A \longrightarrow A \otimes_R A \longrightarrow A \quad (4)$$

of Proposition 5.1.9 of [4] is exact (where we define $A := A_F$), if and only if the complexes of (1) are quasi-isomorphic to S_i , for each i . It is known that the sequence (4) is exact at every place, except possibly at the first two nonzero places. The injectivity of the map j is proved in [6], using consistency. From the gradings we have endowed on the bimodules E , E^* and L , this is a complex of G_1 -graded A -bimodules. Denote the complex (4) by $L^\bullet$. Since there is only one position in which $L^\bullet$ may have nontrivial homology, we deduce that

$$\sum_{n \in \mathbb{Z}} (-1)^n \dim(H^n(L^\bullet)_\nu) = 0$$

for all ν morphisms in the groupoid G_1 , if and only if $L^\bullet$ is exact. This is equivalent to

$$\sum_{n \in \mathbb{Z}} (-1)^n \dim((L^n)_\nu) = 0$$

for all ν morphisms in the groupoid G_1 . For each vertex i we have a complex $M_i^\bullet$ of right A -modules given by the natural morphism from the complex of (1) to the simple module S_i . $M_i^\bullet$ is exact if and only if the complex (1) is a projective resolution of S_i . Let $[u]$ be a G_1 -grade corresponding to some path u . We have seen, in the proof of Theorem (4.1), that $(M_i^\bullet)_{[u]}$ is exact if u is not minimal. Furthermore, in the case in which u is minimal, we showed that

$$\sum_{n \in \mathbb{Z}} (-1)^n \dim(M^n)_{[u]} = 1 \neq 0$$

if $(M^n)_{[u]}$ is not exact: this is situation (B) of that proof. In particular, $M^\bullet$ can only have nontrivial homology in one position.

Given a right A_F -module M we can consider M as a right R -module via the inclusion map $\iota : R \rightarrow A_F$. If M is a right A_F -module of the form $[u]A$, we give M a *left* R -action by setting

$$\begin{aligned} 1_i \cdot m &= m & \text{if } i = t(u) \\ &= 0 & \text{otherwise.} \end{aligned}$$

Then each $M_i^\bullet$ is a complex of R -bimodules. Set

$$M^\bullet := \bigoplus_{i \in V(Q)} M_i^\bullet.$$

This, again, is a G_1 -graded complex of R -bimodules. Now observe that the objects in the complex $A_F \otimes_R M^\bullet$ are the same as those of the complex $L^\bullet$ (though the morphisms are different). By our criterion for (left) faithful flatness of an R -bimodule, we deduce that $A_F \otimes_R M^\bullet$ is exact if and only if all of the $M_i^\bullet$ are, and also $A_F \otimes_R M^\bullet$ can only have nontrivial homology in one position, since $M^\bullet$ can only have nontrivial homology in one position. It follows that

$$\sum_{n \in \mathbb{Z}} (-1)^n \dim(H^n(L^\bullet)_{[u]}) = \sum_{n \in \mathbb{Z}} (-1)^n \dim((L^n)_{[u]})$$

is zero if and only if $(M_{t(u)}^\bullet)_{[u]}$ is exact. The theorem follows. $\square$

Corollary 4.4. *If A_F is the algebra associated to a consistent brane tiling F on a surface M_g , with $g \geq 1$, then A_F is Calabi-Yau of dimension 3.*

The next theorem is stated without proof. The proof uses only those ideas found in the proof of Theorem (4.1). A winding argument is again used.

Theorem 4.5. *If $A = A_F$ is the algebra associated to a consistent tiling F , there is an injective resolution of A as a left A -module:*

$$\begin{aligned} A \xrightarrow{ev} \bigoplus_{i \in V(Q)} \text{Hom}_{\mathbb{C}}(e_i A, e_i A) &\xrightarrow{d} \bigoplus_{a \in E(Q)} \text{Hom}_{\mathbb{C}}(e_{h(a)} A, e_{t(a)} A) \xrightarrow{r} \\ &\xrightarrow{r} \bigoplus_{a \in E(Q)} \text{Hom}_{\mathbb{C}}(e_{t(a)} A, e_{h(a)} A) \xrightarrow{p} \bigoplus_{i \in V(Q)} \text{Hom}_{\mathbb{C}}(e_i A, e_i A) \end{aligned}$$

and an analogous injective resolution of right A -modules. Here Q is the dual quiver to F , $V(Q)$ are its vertices, and $E(Q)$ are its edges.

The maps are as follows: $ev(x)$ is the linear map given by right multiplication by x . For an arbitrary $a \in E(Q)$ and an element $\phi \in \bigoplus_{i \in V(Q)} \text{Hom}_{\mathbb{C}}(e_i A, e_i A)$

there are two linear maps $e_{h(a)}A \rightarrow e_{t(a)}A$ given by the two routes in the diagram below:

$$\begin{array}{ccc} e_{h(a)}A & \xrightarrow{a \cdot} & e_{t(a)}A \\ \downarrow \phi_{h(a)} & & \downarrow \phi_{t(a)} \\ e_{h(a)}A & \xrightarrow{a \cdot} & e_{t(a)}A \end{array}$$

we set

$$(d(\phi))_a(v) = a \cdot \phi_{h(a)}(v) - \phi_{t(a)}(a \cdot v).$$

Now let $\psi \in \bigoplus_{a \in E(Q)} \text{Hom}_{\mathbb{C}}(e_{h(a)}A, e_{t(a)}A)$. For a given $a \in E(Q)$, $\omega_{a,B}^{br}$ and $\omega_{a,W}^{br}$ give a composition of the linear maps defined by ψ , which we denote $\alpha_{\psi,a,B}$ and $\alpha_{\psi,a,W}$, in $\bigoplus_{a \in E(Q)} \text{Hom}_{\mathbb{C}}(e_{t(a)}A, e_{h(a)}A)$. Let $r(\psi)_a = \alpha_{\psi,a,B} - \alpha_{\psi,a,W}$.

Finally, given a $\nu \in \bigoplus_{a \in E(Q)} \text{Hom}_{\mathbb{C}}(e_{t(a)}A, e_{h(a)}A)$ and given an $i \in V(Q)$, define E_1 as in Theorem (4.1). We define $p(\nu)_i(v) = \sum_{a \in E_1} \nu_a(av)$. This sequence is begun by looking at the injective resolution of a quiver algebra without relations found in [2].

Corollary 4.6. *If F is a consistent tiling then the associated algebra A_F is Gorenstein in the sense of [8].*

This is especially significant in the genus 1 case, since if F is consistent and A_F is module finite over its centre, and the centre is a Gorenstein Noetherian algebra, we have that for the category of A_F -modules, $(-)[3]$ is a Serre functor relative to $\mathcal{K}_{Z(A_F)}$, the dualizing complex for the centre of A_F (Theorem 7.2.14 of [4]). In particular, by viewing finite-dimensional modules over A_F as compactly supported sheaves on $Z(A_F)$ via the forgetful functor, the isomorphism

$$\mathbf{R}\text{Hom}_{Z(A_F)}(\mathbf{R}\text{Hom}_{A_F}(M, N), \mathcal{K}_Z) \cong \mathbf{R}\text{Hom}_{A_F}(N, M[3]) \quad (5)$$

yields the *functorial* isomorphism $\text{Ext}^i(M, N) \cong \text{Ext}^{3-i}(N, M)'$ noted in [6]. Furthermore, if the centre is a 3 dimensional toric variety, (5) restricts to an equivariant functorial isomorphism on the category of T -graded modules, where T is the subtorus of $(\mathbb{C}^*)^{E(\Gamma)}$ preserving the superpotential relations $\partial/\partial(X_e)W$. Here the left hand side of (5) is taken in the category of T -graded modules over $Z(A_F)$, with its T -graded dualizing complex.

5 Dimer configurations

We expand upon Theorem 5.4 and Corollary 5.5 of [6], concerning the link between height functions and finite-dimensional G_1 -graded cyclic A_F -modules, starting with some definitions.

Definition 5.1. *A dimer configuration for a graph Γ is a subset $D \subset E(\Gamma)$ of the edges of Γ such that for every $v \in V(\Gamma)$, a vertex of Γ , there is a unique $e \in D$ containing v . Two dimer configurations D', D for a graph Γ are said to be asymptotic if the complements $D \setminus D'$ and $D' \setminus D$ are finite.*

Definition 5.2. Let F be a tiling on the surface M_g , $g \geq 1$. A height function for $\tilde{F}$ is a function from the tiles in $\tilde{F}$ to the integers.

Definition 5.3. To an ordered pair (D, D') of asymptotic dimer configurations on the 1-skeleton $\tilde{\Gamma}$ of $\tilde{F}$ we associate the height function $h(D, D')$ in the following way: the set of $e \in E(\tilde{\Gamma})$ contained in exactly one of D or D' forms a set of disjoint loops $H_{D, D'}$, which we direct so that if an edge in $H_{D, D'}$ comes from D we direct it from black to white, and vice versa for D' . In this way we obtain a set of disjoint oriented loops. We then pick a point i_∞ outside all of these loops, and set the height of a tile $i \in \tilde{F}$ to be given by the oriented intersection number of these loops with a path from i_∞ to the interior of i .

Definition 5.4. We define an i -cyclic A_F -module to be a module that is isomorphic to the cokernel of $f : N \rightarrow e_i A_F$, for some N an A_F -module, and f some morphism of A_F -modules. We define a G_1 -graded i -cyclic module to be one that is a cokernel of such a f , where f is a morphism of G_1 -graded modules.

Definition 5.5. Let i be a tile of F . After choosing a lift $i_0 \in \tilde{F}$ of i , a finite-dimensional i -cyclic G_1 -graded A_F -module M gives a height function, denoted $H(M)$, in the following way: M has a natural decomposition

$$M \cong \bigoplus_{j \in \tilde{F}} M_j \quad (6)$$

where M_j is spanned as a vector space by G_1 -homogeneous elements of M whose G_1 -grade is represented by a path starting at i_0 and terminating at j . We define $H(M)(j) = \dim(M_j)$.

Throughout this section we fix, once and for all, a lift i_0 of the tile i .

Lemma 5.6. Let F be a consistent tiling. Let M be a G_1 -graded i -cyclic A_F -module. Then M is determined up to isomorphism by $H(M)$.

Proof. Let M be a G_1 -graded i -cyclic A_F -module. Let j_0 be some tile of $\tilde{F}$. Let M_{j_0} be a vector space summand of M , as in the decomposition (6). M_{j_0} carries a G_1 -grading. Let j be the projection of j_0 onto F . We denote by $\mathbb{C}[\omega_j]$ the subalgebra of A_F given, as a vector space, by linear sums of closed j -loops. By consistency, and Lemma (2.8), we have

$$\mathbb{C}[\omega_j] \cong \mathbb{C}[x].$$

This algebra carries a natural G_1 -grading. M_{j_0} is also naturally a G_1 -graded $\mathbb{C}[\omega_j]$ -module.

Next we consider $e_i A_F$ as an i -cyclic A_F -module. It has a decomposition as in (6), and we denote the piece corresponding to the tile j_0 $e_i A_{F, j_0}$. M_{j_0} is a quotient module of $e_i A_{F, j_0}$. The module $e_i A_{F, j_0}$ also has the structure of a $\mathbb{C}[\omega_j]$ -module, and from Lemma (2.8) we deduce that in fact this module is free of rank one. It follows that there is a unique G_1 -graded quotient module of $e_i A_{F, j_0}$ of dimension $H(M)(j_0)$, and so we have determined M_{j_0} . Since j_0 was arbitrary we are done. $\square$

The proof of the following theorem occupies the rest of the section.

Theorem 5.7. *For every consistent brane tiling F on a surface of genus at least 1, and every choice of a tile $i_0 \in \tilde{F}$, a lift of $i \in F$, there is a unique dimer configuration D_{F,i_0} for the 1-skeleton $\tilde{\Gamma}$ of $\tilde{F}$, such that there is a 1-1 correspondence between finite-dimensional i -cyclic G_1 -graded A_F -modules M and dimer configurations for $\tilde{\Gamma}$ that are asymptotic to D_{F,i_0} . If D is a dimer configuration for the 1-skeleton $\tilde{\Gamma}$ of $\tilde{F}$ which is asymptotic to D_{F,i_0} , the corresponding module is given by taking the unique i -cyclic A_F -module M satisfying*

$$H(M) = h(D, D_{F,i_0}).$$

This is essentially a higher genus version of Theorem 5.4 from [6], but as ever we dispense with their second consistency condition and the nondegeneracy condition, since we are using the G_1 -grading and not the torus grading constructed there. We will first prove a number of lemmas regarding height functions associated to i -cyclic G_1 -graded A_F -modules.

An i -cyclic G_1 -graded finite-dimensional A_F -module M is equivalent to a finite collection Ω_M of F -term equivalence classes in $\tilde{F}$ starting at i_0 , satisfying the condition that if $[u]$ is in Ω_M , then $[u'] \in \Omega_M$. To such a module we have already associated the height function $H(M)$. For j an arbitrary tile of $\tilde{F}$, $H(M)(j)$ is the number of equivalence classes of paths in Ω_M that end at j .

Recall the construction of $h(D_1, D_2)$ for two arbitrary asymptotic dimer configurations: we first obtain from the ordered pair (D_1, D_2) a disjoint union of oriented loops, and then to a disjoint union of oriented loops we associate a height function. If all the loops are oriented the same way, this height function, applied to some tile in $\tilde{F}$, just counts the minimum number of loops one must cross to get from the interior of the tile to infinity.

Lemma 5.8. *For all G_1 -graded i -cyclic finite-dimensional A_F -modules M , $H(M)$ is given by disjoint loops.*

Proof. First note that this is really a local statement. If we define $\mathcal{T}_V$ to be the set of tiles containing some vertex V of $\tilde{F}$, then the lemma states that the height function, when restricted to $\mathcal{T}_V$, is either constant, or takes precisely two values n and $n+1$, with a line through $\mathcal{T}_V$ composed of two of the edges of F such that on one side of the line $H(M)$ takes value n , and on the other side $H(M)$ takes value $n+1$. If this local statement is true everywhere, the global statement of the lemma easily follows.

We show that $<_i$ restricts to a total order on $\mathcal{T}_V$ (see Definition (2.11) for the definition of $<_i$). Let $j, k \in \mathcal{T}_V$. Let v_{ij} and v_{ik} be minimal paths to the two tiles from i_0 . There is a minimal path from j to k consisting of arrows going around the vertex V , which we denote v_{jk} . We define v_{kj} similarly. Clearly we cannot have that $v_{ij}v_{jk}$ and $v_{ik}v_{kj}$ are both minimal, by antisymmetry of $<_i$. So say, without loss of generality, that $v_{ik}v_{kj} \sim v_{ij}\omega_j^t$, for $t \geq 1$. We have the further decomposition $v_{ij}\omega_j^t \sim v_{ij}\omega_j^{t-1}v_{jk}v_{kj}$, from which we deduce, by

consistency, that $v_{ik} \sim v_{ij}\omega_j^{t-1}v_{jk}$. By minimality of v_{ik} we deduce that $t = 1$, and that $v_{ij}v_{jk}$ is minimal. We deduce that $j <_i k$; thus we have proved that $<_i$ induces a total order on $\mathcal{T}_V$.

Next note that if $j <_i k$ then $H(M)(j) \geq H(M)(k)$, since by the assumption $j <_i k$ we have a minimal path v_{ik} with subpath v_{ij} a minimal path to j , and

$$\{[v_{ik}], [\omega_i v_{ik}], \dots, [\omega_i^{H(M)(k)-1} v_{ik}]\} \subset \Omega_M,$$

and so

$$\{[v_{ij}], [\omega_i v_{ij}], \dots, [\omega_i^{H(M)(k)-1} v_{ij}]\} \subset \Omega_M.$$

Similarly, if j and k share a vertex V then the difference between $H(M)(j)$ and $H(M)(k)$ is at most 1, since if $H(M)(j) = t$ then

$$\{[v_{ij}], [v_{ij}\omega_j], \dots, [v_{ij}\omega_j^{t-1}]\} \subset \Omega_M$$

and so

$$\{[v_{ij}v_{jk}], [v_{ij}\omega_j v_{jk}], \dots, [v_{ij}\omega_j^{t-2} v_{jk}]\} \subset \Omega_M.$$

We have shown that $<_i$ induces a total order of $\mathcal{T}_V$. Let k_1 be the maximum with respect to this ordering. Let k_2 be the maximum element of $\mathcal{T}_V \setminus k_1$, with respect to $<_i$. Then there is a minimal path u such that u decomposes as $u = vw$, where v is a minimal path from i to k_2 and w is a minimal path from k_2 to k_1 . Since k_1 and k_2 share a vertex V , there is a minimal path from k_2 to k_1 only crossing those edges of $\tilde{F}$ that contain V . It follows that the path w consists of a solitary arrow, for else there is some $k_3 \neq k_1$ such that k_3 contains V , and such that $k_2 <_i k_3$. We deduce that, assuming the dual quiver Q is oriented clockwise around V (i.e. V is coloured black), the tiles are arranged, in order with respect to $<_i$, clockwise. If, on the other hand, V is coloured white, then the tiles are arranged, in order with respect to $<_i$, anticlockwise.

The function $H(M)$ is monotone decreasing with respect to the ordering $<_i$, and the maximum difference between $H(M)(j)$ and $H(M)(k)$ for $k, j \in \mathcal{T}_V$ is 1. It follows that, at V , the height function is given by a single line through V , passing along two of the edges of $\tilde{F}$ containing V , or by no lines at all. This implies that the height function is given by a disjoint union of loops. $\square$

Now, let $H(M)(i) = n$, then consider the G_1 -graded short exact sequence

$$0 \longrightarrow [\omega_i]^{n-1}M \longrightarrow M \longrightarrow N \longrightarrow 0.$$

The height function $H([\omega_i]^{n-1}M)$ has maximum value 1, since it takes value 1 at i_0 . If a G_1 -graded i -cyclic module M satisfies

$$\text{image}(H(M)) \subset [0, 1]$$

we call M a *bungalow*. An i -cyclic G_1 -graded submodule M' of M is determined by the minimum value of r such that $[\omega_i]^r \in M'$ (identifying M with a quotient




Figure 3: winding around the loop $\mathcal{L}$

of $e_i A_F$). If M' is a bungalow this value is forced, and must be $n - 1$. We deduce that $[\omega_i]^{n-1} M$ is the unique sub-module of M that is a bungalow. If

$$M' \rightarrow M \rightarrow M''$$

is a G_1 -graded short exact sequence it follows from the construction of the height functions associated to the three modules that $h(M) = h(M') + h(M'')$. Note that if M is i -cyclic, M'' is too. We deduce, by induction on $h(M)$, that a G_1 -graded finite-dimensional cyclic module has a unique filtration by bungalows. We will need the following lemma:

Lemma 5.9. *The height function $H(M)$ associated to a bungalow M is given by a single loop.*

Proof. Let H_1 be the set of tiles in $\tilde{F}$ sent to 1 by $H(M)$. H_1 is connected, since $H(M)$ is monotone decreasing with respect to $<_i$. We need to show that it is simply connected. Unsurprisingly, this will involve winding. Let $\mathcal{L}$ be a loop of edges of $\tilde{\Gamma}$ bounding a hole in H_1 . Since we are working on a universal cover of a surface of genus at least 1, it makes sense to talk of the inside and the outside of this loop - we let the interior side be the side containing none of the tiles of H_1 . Since $H(M)$ is the height function associated to a module, it follows by Lemma (5.8) that it is given by disjoint loops, and so all of the tiles outside $\mathcal{L}$ with boundary intersecting $\mathcal{L}$ nontrivially are contained in H_1 . Let a, b, c be

three arrows as indicated in Figure (3) (we label the edges of Γ dual to these arrows). Let u_0 be a minimal path to $h(a)$. There is a minimal path v_0 from $h(a)$ to $t(b)$ as indicated in Figure (3). We claim that u_0v_0 is minimal. To see this, assume otherwise. Then $u_0v_0 \sim w\omega_{t(b)}$ for some path w . we can pick $\omega_{t(b)}$ so that it decomposes as w_0v_0 , where w_0 is as in Figure (3). By consistency it follows that $u_0 \sim ww_0$. since $H(M)$ is monotone decreasing with respect to $<_i$ we deduce that ww_0 passes only through tiles contained in H_1 , but we picked w_0 to leave H_1 , a contradiction. we deduce that u_0v_0 is minimal.

Let u_1 be a minimal path to $h(c)$. By exactly the same argument, if v_1 is the minimal path from $h(c)$ to $t(b)$ indicated in Figure (3) then u_1v_1 is minimal. By consistency, all minimal paths are equivalent, and so there is a chain of basic F-term equivalent paths taking us from u_0v_0 to u_1v_1 . These paths must all be contained in H_1 , since $H(M)$ is monotone decreasing with respect to $<_i$. Let T be a tubular neighborhood of $\mathcal{L}$ (part of T is indicated in Figure (3)). Let γ_i satisfy $\Psi(\gamma_i) = u_i$ for $i = 0, 1$. Let λ_i satisfy $\Psi(\lambda_i) = v_i$ for $i = 0, 1$. As indicated, we can choose λ_i to lie entirely within T , and for γ_i to end at the starting point of λ_i , for each i . Then we can construct a homotopy from γ_0 to γ_1 such that $\text{image}(H|_{[0,1] \times \{1\}}) \subset T$, by the following procedure. First, take a homotopy that stretches γ_0 to the path given by concatenating γ_0 and λ_0 . Similarly, there is a homotopy that stretches γ_1 along the path λ_1 marked in Figure (3). Finally, by F-term equivalence of u_0v_0 and u_1v_1 , there is a homotopy between these two stretched paths. We can repeat this argument for the next two edges of $\mathcal{L}$, proceeding clockwise around the loop, until we arrive back at a . We can pick T to be arbitrarily thin. We have shown that there is a γ_0 such that $\Psi(\gamma_0) = u_0$, and a γ_1 such that $\Psi(\gamma_1) = u_1$, and a homotopy

$$H : [0, 1]^2 \rightarrow H_1$$

such that $H|_{\{i\} \times [0,1]} = \gamma_i$, and $\text{im}(H|_{[0,1] \times \{1\}}) \subset T$. After repeatedly applying this argument we arrive at a γ_∞ such that $\Psi(\gamma_\infty) \sim u$, but γ_∞ doesn't have the same winding number around $\mathcal{L}$ as γ_0 , a contradiction since γ_0, γ_∞ are k -unwound for any tile k not in H_1 . This proves the lemma. $\square$

So far we have proved that an i -cyclic G_1 -graded module M is determined by its filtration by bungalows. Furthermore, a bungalow determines a single loop in $\tilde{F}$, defined by its height function. The height function for M is given by the disjoint union of these loops, and M is uniquely determined by this disjoint union of loops. In one direction, we have shown that a module determines a finite disjoint union of loops that satisfy the property that they are the height functions of bungalows. By the next lemma, this is a 1-1 correspondence

Lemma 5.10. *If H is the height function associated to a finite disjoint union of loops arising from bungalows, then H is the height function associated to a G_1 -graded i -cyclic module.*

Proof. Let $\mathcal{L}_s$ $s \in [1, \dots, n]$ be a finite union of disjoint loops, each the height function of a bungalow M_s . Since they are disjoint, and all contain i , they form

a series of concentric loops around i , ordered by inclusion. We label them so that $\mathcal{L}_s$ contains $\mathcal{L}_t$ if $s < t$. For j a tile of $\tilde{F}$, let v_{ij} denote a minimal path from i_0 to j (these are all equivalent, by consistency). Let

$$\Omega_s := \{[v_{ij}][\omega_j]^r | j \in \tilde{F}, r \geq s \text{ if } j \text{ is inside } \mathcal{L}_s, r \geq (s-1) \text{ otherwise}\}$$

Now a path that goes outside $\mathcal{L}_s$ and then back inside cannot be minimal, since $H(M_s)$ is monotone decreasing with respect to $<_i$. It follows that if u is a path such that $[u] = [v_{ij}][\omega_j]^{s-1}$, and j is outside $\mathcal{L}_s$, then if v is a path from j to k , for k inside $\mathcal{L}_s$, we must have $[uv] = [v_{ih(k)}][\omega_j]^t$ for $t \geq s$. We deduce that Ω_s is a basis for a sub- A_F -module of A_F (i.e. a right ideal), which we denote N_s . Now, by construction, the cokernel of the morphism

$$\bigoplus_{s \in [1, \dots, n]} N_s \rightarrow A_F$$

is an i -cyclic A_F -module with the required height function. $\square$

Let D be a dimer configuration on $\tilde{F}$. We say that a loop $\mathcal{L}$ is *obtainable* from D if alternating edges of $\mathcal{L}$ belong to D . It is clear enough that the set of dimer configurations D' on $\tilde{F}$ that are asymptotic to D is in 1-1 correspondence with the set of sets of disjoint obtainable loops in $\tilde{F}$. This correspondence is realised via the construction $h(D, D')$. We are finally in a position to prove Theorem (5.7), since we have reduced it to the following Proposition:

Proposition 5.11. *For every consistent brane tiling F on a surface of genus at least 1, and every choice of a tile $i_0 \in \tilde{F}$, there is a unique dimer configuration D_{F, i_0} for the 1-skeleton $\tilde{\Gamma}$ of $\tilde{F}$, such that the obtainable loops of D_{F, i_0} are exactly the loops corresponding to height functions of bungalows.*

Proof. We extend the partial order $<_i$ to a total order $<$, and obtain in this way a function κ from the tiles in $\tilde{F}$ to $\mathbb{N}$. Let $\mathfrak{T}_n = \kappa^{-1}([1, \dots, n])$. We assume that we have constructed a dimer configuration D_n on $\mathfrak{T}_n$ satisfying the condition above - the set of loops in the 1-skeleton of $\mathfrak{T}_n$ with alternating edges in D_n is in bijective correspondence with the set of loops in the 1-skeleton of $\mathfrak{T}_n$ that arise from bungalows. For $n = 1$, existence and uniqueness of this configuration is clear.

Note that the boundary of $\mathfrak{T}_n$ is the loop corresponding to the bungalow which is the cokernel of the map

$$[\omega_i]A \oplus \bigoplus_{j \notin \mathfrak{T}_n} [v_{ij}]A \longrightarrow e_i A$$

and so the correct alternating edges of the boundary of $\mathfrak{T}_n$ belong to D_n , by induction. Figure (4) depicts a part of the loop bounding $\mathfrak{T}_n$, along with the tile $\kappa^{-1}(n+1)$. Recall from the proof of Lemma (5.8) that for each vertex j , the restriction of $<_i$ is already a total order, and around the white vertices the tiles are arranged anticlockwise, with respect to this ordering, and around the black

vertices they are arranged clockwise. It follows that the addition of the new tile must take the form as depicted in Figure (4), with the part of the loop bounding $\mathfrak{T}_{n+1}$ that is not contained in $\mathfrak{T}_n$ being a path which, travelling clockwise, goes from a white to a black vertex. Since we require the boundary of $\mathfrak{T}_{n+1}$ to be a loop obtainable from D_{n+1} , from Figure (4) it is clear how we *must* (hence, uniqueness) extend D_n , along the edges marked in bold, where the edges that must already be in D_n are also marked in bold. We now give the inductive step: assume we have a loop S , obtainable from D_{n+1} , contained in the 1-skeleton of $\mathfrak{T}_{n+1}$. We assume that it encloses $\kappa^{-1}(n+1)$. Then from the figure if we modify the loop by removing $\kappa^{-1}(n+1)$ from its interior, we obtain a loop in the 1-skeleton of $\mathfrak{T}_n$ obtainable from D_n . By induction this loop S' corresponds to a bungalow given by the cokernel of the map

$$[\omega_i] \oplus \bigoplus_{j \notin \text{Int}(S')} [v_{ij}] A \longrightarrow e_i A.$$

Any minimal path to $\kappa^{-1}(n+1)$ must pass through one of the tiles in $\mathfrak{T}_n$ adjoining $\kappa^{-1}(n+1)$, and so must lie entirely within S' , until it leaves for $\kappa^{-1}(n+1)$. It follows that the set P of paths $u \in \text{paths}_F$ that are minimal paths from i_0 to tiles in the interior of S is closed under the operation $u \rightarrow u'$ of deleting the final arrow, and so if Q is the set of F-term equivalence classes of paths not in P , Q spans a right ideal of A_F , which we call I_Q . By construction, the cokernel of

$$I_Q \longrightarrow e_i A$$

gives the required module. Conversely, say we have a bungalow M with associated loop S contained in $\mathfrak{T}_{n+1}$. Without loss of generality it encloses $\kappa^{-1}(n+1)$, and we set $j = \kappa^{-1}(n+1)$. Then the bungalow given by the cokernel of the map


$$[v_{ij}] A \longrightarrow M$$

gives a loop contained in $\mathfrak{T}_n$, and, by induction, alternating edges are contained in D_n . We deduce from the indicated extension of D_n that the loop S has alternating edges in D_{n+1} . This completes the proof of the Proposition, and the proof of Theorem (5.7). $\square$

6 Donaldson-Thomas theory

The final section of [6] deals with the Donaldson-Thomas theory of the moduli space $\mathcal{M}_{\mathbf{v},i}$ constructed in [7], of framed i -cyclic modules with dimension vector $\mathbf{v} \in \mathbb{N}^{V(Q)}$. We consider the moduli of pairs (M, f) , where M is an i -cyclic A_D -module, and $f : e_i A_F \rightarrow M$ is a surjection of A_F -modules. Given an action of some torus T on the moduli space, such that the perfect obstruction theory on the moduli space is equivariant with respect to T , and the fixed locus consists of isolated fixed points, Corollary 3.5 of [5] states that the Donaldson-Thomas invariant of the moduli space is given by the sum, over the T -fixed points, of the

Figure 4: The bold edges show those from D_n , as well as new ones for D_{n+1} .



sign of the dimension of the Zariski tangent space at those points. In order to apply Corollary 3.5 of [5] we need to find the fixed points of the moduli space, under some action of a torus T that respects the perfect obstruction theory, confirm that the Zariski tangent space has no trivial weights under the induced action of T , and then compute the dimension of the Zariski tangent spaces.

The torus $(\mathbb{C}^*)^{|E(F)|}$ acts naturally on the algebra A_F . If $T' \subset (\mathbb{C}^*)^{|E(F)|}$ is the subtorus preserving the superpotential relations, then there is a natural action of T' on $\mathcal{M}_{\mathbf{v},i}$. This action is not faithful, and the kernel of the map

$$\theta : T' \rightarrow \prod_{\mathbf{v} \in \mathbb{N}^V(Q)} \text{Aut}(\mathcal{M}_{\mathbf{v},i}) \quad (7)$$

is given by the subtorus generated by those 1-parameter subgroups of T' that rescale all arrows going into one vertex by some number $z \in \mathbb{C}$, and rescale all arrows going out by z^{-1} . We let T be the image of θ (again T is a torus). Let T_W be the image under θ of the largest subtorus of T' acting with trivial weight on the simple loop ω_i . The obstruction theory constructed in [7] is symmetric equivariant with respect to the induced action of T_W .

Lemma 6.1. *The torus T_W has dimension $2g$.*

Proof. We show the coweight lattice of T_W is $\mathbb{Z}^{2g}$. Pick i a tile of F , and let $i_0 \in \tilde{F}$ be an i -tile. Let

$$\alpha : \mathbb{C}^* \rightarrow T_W$$

be a coweight. α is determined by a $\mathbb{C}^*$ -action on A_F that acts trivially on ω_i . Note that by transport of simple loops (see Remark (2.2)) this implies $\mathbb{C}^*$ acts trivially on all simple loops. Such an action is determined by its effect on minimal paths. Since we have picked an initial i_0 , it makes sense to talk of the action of $\mathbb{C}^*$ on a tile in $\tilde{F}$ under α , and in fact α is determined by the actions of $\mathbb{C}^*$ on the tiles of $\tilde{F}$. For any u a j -loop, there is a v which is an i -loop, and a path

w from i to j , such that $wu\omega_j^n \sim vw$ for some $n \geq 0$. It follows that the $\mathbb{C}^*$ -action on j -loops is determined by the $\mathbb{C}^*$ -action on i -loops. Furthermore, increasing the weights on all j -tiles corresponds to an action of the gauge group. It follows that the action of $\mathbb{C}^*$ on $\prod_{\mathbf{v} \in \mathbb{N}^{V(Q)}} \text{Aut}(\mathcal{M}_{\mathbf{v},i})$ determines and is determined by its action on the i -tiles of $\tilde{F}$. Such an action uniquely determines an element of $H_1(M_g, \mathbb{Z})' \cong \mathbb{Z}^{2g}$, and so all that is required is to show that every element of $H_1(M_g, \mathbb{Z})'$ is realised in this way.

Assume that we have drawn the dual graph of F on M_g , so we have some embedding $Q \subset M_g$. Let C be a curve in M_g representing some $\beta \in H_1(M_g, \mathbb{Z})$. We can pick C so that it doesn't go through the vertices of Q . Let $p^{-1}(C)$ be the preimage of C under the projection from the universal cover. Now number the tiles in $\tilde{F}$ so that the difference between two adjacent tiles j and k is given by the oriented intersection number of the arrow between the vertices in the dual graph corresponding to j and k , with $p^{-1}(C)$. This gives an assignment of numbers to the tiles of $\tilde{F}$. By construction, the corresponding element of $H_1(M_g)'$ is the image of β under the natural isomorphism induced by the nondegenerate intersection product. This completes the proof. $\square$

Remark 6.2. Define T'_W to be the largest sub-torus of $(\mathbb{C}^*)^{|E|}$ preserving the superpotential relations and acting with trivial weight on the simple loop ω_i . In the course of the above proof we have shown that T'_W acts with trivial weight on all simple loops.

Proposition 6.3. If F is a consistent tiling on a surface of genus 1, then for all $\mathbf{v} \in \mathbb{N}^{V(Q)}$ the T_W -fixed points of $M_{\mathbf{v},i}$ are the G_1 -graded quotients of $e_i A$ with dimension vector $\mathbf{v}$.

Remark 6.4. In fact the above holds if and only if F is consistent.

Proof. Let M be a framed i -cyclic A_F -module, and let

$$I \rightarrow A_F \rightarrow M$$

be the associated short exact sequence. We consider I as a right ideal of A_F . I is generated by elements of the form

$$\sum_{s \in S} \lambda_s [u_s], \quad (8)$$

where the $\lambda_s \in \mathbb{C}$ and u_s are paths from i to some fixed $j \in F$. Since M is T_W invariant, I is generated by elements of the form in (8), chosen such that the lifts of the u_s to paths starting at i_0 in $\tilde{F}$ all terminate at the same lift j_0 of j . We deduce that, as a vector space, we have a decomposition

$$N \cong \bigoplus_{k_0 \in \tilde{F}} I_{k_0} \quad (9)$$

where each I_{k_0} is a sub- $\mathbb{C}[\omega_{k_0}] (\cong \mathbb{C}[x])$ -module of $e_i A_{F,k_0}$, where $e_i A_{F,k_0}$ is the summand associated to the tile k_0 in the analogous decomposition of $e_i A_F$.

By consistency, each $e_i A_{F,k_0}$ is isomorphic to the free $\mathbb{C}[x]$ -module, and its inherited G_1 -grading is just the usual $\mathbb{N}$ -grading on $\mathbb{C}[x]$, where $\mathbb{N}$ is considered as a category with a unique morphism from n to m if $m \geq n$, and we embed this category into G_1 by the functor which sends n to $u\omega_{h(u)}^n$, and the morphism from n to m to the path $\omega_{h(u)}^{m-n}$, where u is a minimal path from i_0 to k_0 . All that is required is to show that in fact each I_{k_0} is a $\mathbb{N}$ -graded submodule of $e_i A_{F,k_0}$.

We proceed by induction on the codimension of I_{k_0} in $e_i A_{F,k_0}$. If it is zero, there is nothing to prove. Assume it is n , and we have proved the result for all k_0 such that the codimension of I_{k_0} is less than n . Let u be a minimal path from i_0 to k_0 . By Lemma (7.3) of the next section, we deduce that there is an arrow a_0 such that ua_0 is minimal. Denote $h(a_0)$ by k_1 . The ideal $I_{k_0} \subset e_i A_{F,k_0}$ is generated by some degree n polynomial $p(x)$ (in ω_{k_0}), i.e. $up(\omega_{k_0}) \in N$. Since I is a right ideal, we deduce that $ua_0p(\omega_{k_1}) \in I$, and the same polynomial belongs to I_{k_1} . We deduce that the codimension of I_{k_1} is less than or equal to n . If it is equal to n , then in fact right multiplication by a_0 induces an isomorphism from I_{k_0} to I_{k_1} , since any other degree n polynomial contained in I_{k_1} must be a multiple of p . We obtain in this way a chain of isomorphisms until, by finiteness and the fact that minimal paths do not revisit tiles, we arrive at some k_t , and arrow a_t to some k_{t+1} , such that the codimension of I_{k_t} is strictly greater than the codimension of $I_{k_{t+1}}$. $I_{k_{t+1}}$ is an ideal in $e_i A_{F,k_{t+1}}$, generated by some degree l polynomial q . By induction, this polynomial is $q(x) = x^l$. Right multiplication by $\omega_{a_{t+1}}^{br}$ defines a $\mathbb{C}[x]$ -module map from $I_{k_{t+1}}$ to I_{k_t} , which takes $ua_0, \dots, a_t\omega_{k_{t+1}}^l$ to $ua_0, \dots, a_{t-1}\omega_{k_t}^{l+1}$. Since we choose $ua_0, \dots, a_{t-1}$ to be minimal, this represents the element x^{l+1} under the isomorphism $I_{k_t} \cong \mathbb{C}[x]$. Since I_{k_t} has codimension n in $e_i A_{F,k_t}$, it follows that $l = n - 1$ and $p(x)$ is a multiple of x^n . $\square$

We next prove the following:

Proposition 6.5. *If F is a tiling on a surface of genus 1, and $M_{\mathbf{v},i}$ is the moduli space associated to the dimension vector $\mathbf{v}$ then the action of T_W on the Zariski tangent space at the fixed points carries no trivial representations.*

Proof. Let I_0 be the kernel of the map $f : e_i A \rightarrow M$. We consider I_0 as a right ideal of A_F . As in Lemma 4.1(b) of [5], we check the condition via the identification of the Zariski tangent space with the space $\text{Hom}(I_0, M)$. Note that both I_0 and M are G_1 -graded by Proposition (6.3).

Let $T'_W \subset T'$ be the subtorus of $(\mathbb{C}^*)^{|E|}$ preserving the superpotential relations and acting with trivial weight on the simple loops. $\text{Hom}(I_0, M)$ carries a natural T'_W -action, given by the action of T'_W on the tangent space of a fixed point. This coincides with the natural T'_W -action given on homomorphisms between modules equipped with a T'_W -action. The T'_W -trivial sub-representations are just the T_W -trivial sub-representations, by the definition of T_W .

Any A_F -module homomorphism from I_0 to M takes paths ending at j -tiles to paths ending at j -tiles. A homomorphism is acted on trivially by T_W (and

hence by T'_W) if and only if it takes elements of the form $[v_{ij_0} \omega_{j_0}^n]$, where v_{ij_0} is some minimal path from i_0 to j_0 in $\tilde{F}$, to linear combinations of elements of the form $[v_{ij_0} \omega_{j_0}^m]$, with the same $j_0 \in \tilde{F}$. Now let $\phi \in \text{Hom}(I_0, M)$ be a nonzero homomorphism acted on trivially by T_W .

By Proposition (6.3), M and I_0 are G_1 -graded. Recall from the proof of Theorem (5.7) that the module M is equivalent to the data of its height function, and this is given by a set of disjoint loops, each one the height function of one of the modules occuring in its filtration by ‘bungalows’. These loops form a concentric series of loops around i , since they are disjoint and all contain i . Let the loops be ordered so that the outermost is $\mathcal{L}_1$, i.e. $\mathcal{L}_t$ encloses exactly those tiles j such that $H(M)(j) \geq t$. Let j lie outside $\mathcal{L}_1$. Since ϕ is acted on trivially by T_W it preserves the endpoints of paths. It follows that $\phi([v_{ij}]) = 0$, where v_{ij} is the minimal path from i to j , since the G_1 -graded pieces of M corresponding to paths ending at j are all zero. Similarly, for any $[v] \in I_0$ with $h(v) = j$, $\phi([v]) = 0$.

Say, by induction, that for all j outside $\mathcal{L}_t$ it is shown that, for all $[v] \in I_0$ with $h(v) = j$, $\phi([v]) = 0$. Let j be a tile on the interior of $\mathcal{L}_t$, sharing an edge with $\mathcal{L}_t$. Let k be the other tile sharing this edge (note that k lies outside $\mathcal{L}_t$, and $H(M)(k) = t - 1$). Then the two minimal paths v_{kj} and v_{jk} satisfy $v_{jk}v_{kj} \sim \omega_j$. By the definition of $\mathcal{L}_t$, $[v_{ij}][\omega_j^{t-1}][v_{jk}] \in I_0$, and by induction $\phi([v_{ij}][\omega_j^{t-1}][v_{jk}]) = 0$. Any $[v] \in I_0$ with $h(v) = j$ can be written $[v_{ij}][\omega_j^{t-1}][v_{jk}][v_{kj}][\omega_j^r]$, for some $r \geq 0$. It follows that for all $[v] \in I_0$ with $h(v) = j$, $\phi([v]) = 0$.

Now let j and k be two arbitrary neighbouring tiles, both lying inside $\mathcal{L}_t$ and outside $\mathcal{L}_{t+1}$. Let a be the arrow between them, without loss of generality it goes from j to k . Either $v_{ij}a$ is minimal, or $v_{ik}\omega_a^{br}$ is. Consider the decomposition

$$M \cong \bigoplus_{l \in \tilde{F}} M_l$$

where M_l is spanned by those G_1 -graded pieces of M corresponding to paths that end at l . If $v_{ij}a$ is minimal, then right multiplication by a defines an isomorphism from M_j to M_k . Similarly, if $v_{ik}\omega_a^{br}$ is minimal then right multiplication by ω_a^{br} defines an isomorphism from M_k to M_j . Either way, we deduce that $\phi([v_{ij}\omega_j^t]) = 0$ if and only if $\phi([v_{ik}\omega_k^t]) = 0$. This completes the inductive step, and the proposition follows. $\square$

Given these two propositions, we can compute Donaldson-Thomas invariants, assuming only consistency, thanks to the characterisation of the G_1 -graded cyclic modules of Theorem (5.7). We have proved all the essential results of [6] assuming only consistency.

7 Other notions of consistency

We recall the definitions found in [6].

Definition 7.1. A brane tiling F satisfies ‘MR2’ if, for any two tiles $i, j \in \tilde{F}$, there exists an arrow a from j to some tile k such that some minimal path from i to k passes along a . We also require the dual to hold.

Definition 7.2. A brane tiling F is nondegenerate if its 1-skeleton Γ satisfies the following condition: for each edge $E \in E(\Gamma)$ there is a dimer configuration for Γ containing E .

The consistency condition of Definition (2.5), and the two conditions above, are assumed in [6], and are used to prove the results there. It is natural to wonder whether they follow from the consistency condition (the ‘first consistency condition’ in [6]). The answer, at least for the second consistency condition, is affirmative.

Lemma 7.3. Consistency implies condition MR2.

Sketch proof: Pick $i, j \in \tilde{F}$ violating the condition. Define E_1 as the set of arrows a with $h(a) = j$, and E_2 as the set of arrows with $t(a) = j$. For each $k, l \in \tilde{F}$ let v_{kl} denote a minimal path from k to l . Then we are assuming that for each $a \in E_2$, $v_{ij}a \sim v_{ih(a)}\omega_{h(a)}^t$ for some $t \geq 1$. If $t > 1$ we pick $\omega_{h(a)}$ going through j and deduce that v_{ij} is not minimal. It follows that for all $a \in E_2$, $v_{ij} \sim v_{i,h(a)}\omega_{a,B}^{br} \sim v_{ih(a)}\omega_{a,W}^{br}$. None of these paths go through j (except to end at it) by minimality of v_{ij} . We are now in exactly the situation of the winding argument of (4.1), giving a contradiction. The dual argument is identical. $\square$

So condition MR2 is redundant, and if we include our consistency condition we can drop it from our definition of a ‘good’ algebra. While it is possible that nondegeneracy follows from consistency too, we have no proof of this (though see [1]).

8 Final remarks

Theorem 8.1. Let F be a consistent brane tiling on an oriented surface M_g . The associated algebra is 3-Calabi-Yau if and only if $g > 0$.

Proof. From Proposition (4.3), and Theorem (4.1), it is enough to show that the complex (1) is not a resolution of S_i in the genus zero case. For this, it is sufficient to show that one of the G_1 -graded pieces corresponding to a path of length at least 1 is not exact. Let F be a consistent brane tiling on S^2 . From the proof of Theorem (4.1) it is enough to show, using the same notation, that there is a pair of tiles i and j such that for u a minimal path between them $E'_1 = E_1$, i.e. that we are in situation (B) of that proof.

In the case of the sphere we have $F = \tilde{F}$, and Lemma (2.8) applies to F . Now for two tiles i and j of F let v_{ij} be a minimal path between them, and let $S(i, j)$ be the set of tiles contained in paths that are F-term equivalent to v_{ij} . Say we have picked i and j so that $S(i, j)$ is maximal. Note that, for $a \in E_1$, $t(a)$ is in $S(i, j)$ if and only if there is a minimal $v \sim v_{ij}$ starting with ω_a^{br} , i.e.

for $a \in E_1$, $a \in S(i, j)$ if and only if $a \in E'_1$. Say there is some $a \notin E'_1$. It follows that av_{ij} is minimal, since if $av_{ij} \sim \omega_{t(a)}v \sim a\omega_a^{br}v$ for some v , we deduce that $a \in E'_1$. But now clearly $|S(t(a), j)| > |S(i, j)|$, a contradiction. $\square$

Remark 8.2. *Another topological argument shows that in fact $S(i, j)$ contains all the tiles of F . Briefly, we use the same argument as above to deduce that all the tiles sharing edges with j are contained in $S(i, j)$. Now, say $k \notin S(i, j)$ for some $k \in F$. For each tile bordering j there is a tile bordering i , with a $w \sim 'v'_{ij}$ going between the two, for some choice of minimal v_{ij} . We wind around j using these paths, necessarily winding around i too as we go ($S^2 \setminus \{i, j\}$ is an annulus). After winding round j we obtain two F -term equivalent paths from a tile bordering j to a tile bordering i , which form a cycle around the hole left by k in $S(i, j) \setminus \{i, j\}$. This proof is suggestive: a consistent tiling on a surface M_g is not Calabi-Yau if and only if we can find the topological obstruction to M_g being a $K(\pi, 1)$ surface (i.e. $\pi_2 \neq 0$), in one of the $S(i, j)$.*

Remark 8.3. *Let F be a consistent brane tiling on M_g , with $g > 1$, and let $i \in F$. The centre of the algebra A_F is isomorphic to a subalgebra of the algebra of i -loops. It follows that, by considering lifts of paths to the universal covering space, we can consider the algebra $\mathbb{C}[\pi_1(M_g)]$ as a $Z(A_F)$ algebra, and there is a surjective $Z(A_F)$ algebra map from the algebra of i -loops to $\mathbb{C}[\pi_1(M_g)]$. It follows that A_F is never module finite over its centre, and there is no chance of it being a noncommutative crepant resolution of its centre. We are also in bad shape for trying to use Proposition 7.2.14 of [4] to establish Serre duality relative to $\mathcal{K}_{Z(A_F)}$, in fact this dualizing complex is clearly concentrated in the wrong degree.*

We finish with some thoughts on the Donaldson-Thomas theory of the ‘higher genus’ 3-Calabi-Yau algebras.

We are fortunate, in the genus 1 case, that the fundamental group of the real torus is Abelian. We can uniquely decompose i -loops (up to F -term equivalence) as $v\omega_i^t$, where v is a minimal path between the endpoints of the loop, and t is some nonnegative integer. After covering $\mathbb{R}^2$ with fundamental domains of the torus, the T_W weight of u is given uniquely by the fundamental domain that $h(u)$ lives in. In the non-Abelian case we are not so fortunate.

Say $h(u)$ is in a fundamental domain corresponding to the element $\alpha \in \pi_1(M_g)$, with $g > 1$. Then any torus weight on u can only be used to tell us the image of α under the natural noninjective map $\pi_1(M_g) \rightarrow H_1(M_g)$, the Abelianization of $\pi_1(M_g)$. The result is that we lose our assurance that the (topological) T_W -fixed locus is the set of G_1 -graded modules, and should expect to see Grassmannians, or worse, in the T_W -fixed locus. Furthermore, our argument for Proposition (6.5) will not do as it stands, and so more work must be done to show that the T_W -fixed subscheme is reduced. We conclude that the Donaldson-Thomas picture for higher genus consistent brane tilings is more complicated, though perhaps more interesting too.

References

- [1] A. Ishii, K. Ueda, *On moduli spaces of quiver representations associated with dimer models*, arXiv:0710.1898 (2007).
- [2] B. Gothen, A. King, *Homological algebra of twisted quiver bundles*, J. London Math. Soc (2005), no. 71(1), 85–99.
- [3] N. Broomhead, *Dimer models and Calabi-Yau algebras*, Ph.D. thesis, 2009.
- [4] V. Ginzburg, *Calabi-Yau algebras*, arxiv:math/0612139.
- [5] K. Behrend, B. Fantechi, *Symmetric obstruction theories and Hilbert schemes of points on threefolds*, Algebra Number Theory (2008), no. 3, 313–345.
- [6] S. Mozgovoy, M. Reineke, *On the noncommutative Donaldson-Thomas invariants arising from brane tilings*, Arxiv:0809.0117.
- [7] B. Szendrői, *Non-commutative Donaldson-Thomas theory and the conifold*, Geom. Topol. **12** (2008), no. 2, 1171–1202.
- [8] A. Yekutieli, *Dualizing complexes, Morita equivalence and the derived Picard group of a ring*, J. London Math. Soc. **60** (1999), no. 3, 723–746.